

SUMMARY INFORMATION - UXOR®

Topical Medicated Gel

HEALTH PROFESSIONAL SUMMARY INFORMATION

This product monograph was prepared based on the approval of Uxor® for sale in Canada and is designed specifically for consumers. For questions about this medication, contact your doctor or pharmacist.

RECOMMENDED USE

Uxor® helps in the prevention of premature ejaculation.

COMPOSITION

Route of administration	Dosage form / dosage	Active ingredients	Non-medicinal Ingredients
Topical	Topical medicated gel. 56.8 g tube.	Benzocaine (4-Aminobenzoic acid, ethyl ester) 7.5%	Glycerin, macrogol 3350, PEG-8 laurate, polyethylene glycol, sodium saccharinate, sorbic acid, water

Reasons for using this product:

Uxor® is a treatment that is applied topically. Its formula is composed of benzocaine (7.5%). The use of Uxor® is supported by clinical studies aimed in treating men who ejaculate prematurely.

The effects of this product:

Uxor® produces a light desensitization of the penis nerves. In doing so, it allows the user to better manage his level of control and delay his ejaculation.

PRECAUTIONS & CONTRAINDICATIONS

Caution(s) and Warning(s):

Premature ejaculation may be due to a condition requiring medical supervision. If this product, used as directed, does not provide relief, discontinue use and consult a health care practitioner.

Contraindication(s):

If the following symptoms appear: weakness, confusion, headache, difficulty breathing and/or pale, gray or blue coloured skin, stop use and consult a health care practitioner. These symptoms may be signs of methemoglobinemia, a rare disorder, which may appear up to 2 hours after use.

Known Adverse Reaction(s):

Hypersensitivity/allergy has been known to occur; in which case, discontinue use.

DRUG INTERACTIONS

There are no known drug interactions between topically applied Uxor[®] and prescribed or non-prescribed drugs. You should contact your healthcare provider if you have questions about your health. If you are taking or thinking of taking prescription drugs, tell your doctor or pharmacist before taking this product.

PROPER USE OF THIS MEDICATION

Usual dose:

Apply +/- 2 grams to glans and dorsal vein of the penis, 5 to 10 minutes before intercourse. Wipe excess gel before proceeding with sexual activity.

Overdose:

If you apply too much product, simply rinse with water. It is important not to apply too much to prevent from transferring the drug into the vagina of your partner. If this occurs, do not attempt to perform sex. Wait until the product flows out of the vaginal cavity. Do not hesitate to contact a health care provider if you experience unusual side effects not listed.

REPORTING UNEXPECTED SIDE EFFECTS

You can report any unexpected adverse reactions (side effects) associated with the use of health products to Orimed's Pharmacovigilance center via one of the following 3 ways:

By email, at pv@orimedpharma.ca

By phone, by calling toll free 1-866-399-9091

By completing a declaration form Canada Vigilance and sending it:

- By fax, toll-free at 1-866-678-6789, or

- By mail to: Canada Vigilance Program

NOTE

For information related to the management of side effects, or regarding your general health, contact your healthcare provider.

STORAGE

Store at room temperature (15-30 ° C).

Keep out of the reach of children. This packaging is not childproof.

FOR MORE INFORMATION

Contact us if you have questions or comments or wish to obtain more information on the product or its use.

SEND AN EMAIL TO:

info@orimedpharma.ca

CALL US:

1.866.399.9091

WRITE US:

Orimed Pharma Inc.

203-1380 Newton Street,

Boucherville (Quebec)
Canada. J4B 5H2

Orimed Pharma Inc., a Division of JAMP Pharma Inc., prepared this document.
Last revised: September 21, 2015.

MO0004
MO0004-E